N-049/80-5-0021
Amprolium and Ethopa bate /Bacitracin methy kne disalicy late /Roxarsone
for Broiker Chickens - W6/CoccI
Amprolium and Ethopa bate/Bacitracin Methylene Disalicylate/Roxarsone
Type B Med. Feed For Broiler Chickens

## Type B Medicated Feed

For use in manufacturing medicated chicken feed

Indications: For increased rate of weight gain and as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from *Eimeria acervulina*, *E. maxima* and *E. brunetti* is likely to occur in broiler chickens raised in floor pens.

## ACTIVE DRUG INGREDIENTS Bacitracin Methylene Disalicylate 333 mg/lb - 2.34 g/lb.\* Roxarsone 2.27 g/lb. **GUARANTEED ANALYSIS** Crude Protein, not less than Lysine, not less than Methionine, not less than Crude fat. not less than Crude fiber, not more than Calcium, not less than. Calcium, not more than..... Phosphorus, not less than Salt<sup>1</sup>, not less than Salt<sup>1</sup>, not more than Sodium<sup>2</sup>, not less than % Sodium<sup>2</sup>, not more than

## **INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials

## MIXING DIRECTIONS

Mix 15 lb. of this Type B medicated feed with 1985 lb. of non-medicated feed to make one ton of Type C medicated feed containing 113.5 g/ton amprolium, 36.3 g/ton ethopabate, 5-35 g/ton bacitracin methylene disalicylate\* and 34 g/ton roxarsone

**CAUTION**: Do not feed to laying chickens. Do not use as a treatment for outbreaks of coccidiosis. Do not use in Type C medicated feeds containing bentonite.

→ WARNING: Withdraw 5 days before slaughter. Use as the sole source of amprolium and organic arsenic. ←

\*The final printed feed label should list only a single drug concentration

Net Weight lb ( kg

Manufactured by BlueBird Feed Mill Anytown, Anystate 12345

<sup>&</sup>lt;sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.